

Hunting related shooting incidents during spring and fall wild turkey seasons, Pennsylvania
 Incident rate is the number of incidents per 100,000 hunters.

YEAR	SPRING			F ^A	
	Nonfatal	Fatal	(HRSIs/100,000 hunters)	Nonfatal	Fatal
1985	8	2	3.7	17	1
1986	10	0	4.1	25	1
1987	8	0	3.9	25	1
1988	8	0	3.5	21	3
1989	4	1	1.8	19	0
1990	8	1	4.2	38	3
1991	16	0	8.9	37	2
1992 ^a	8	0	4.3	6	0
1993 ^b	5	0	2.5	6	0
1994	9	1	4	7	0
1995	16	0	6.7	14	0
1996	12	0	5	11	1
1997	12	0	4.7	14	0
1998	11	0	5.6	8	2
1999	14	0	5.9	16	1
2000	10	0	4.5	10	0
2001	7	0	3	13	0
2002	8	1	3.7	15	0
2003	9	0	3.6	2	0
2004 ^c	9	0	?	5	0
2005	8	0	3.2	2	0
2006	5	0	2.0	4	0
2007	5	0	2.2	1	0
2008 ^d	8	0	3.7	2	0
2009	5	0	2.2	3	0
2010	10	1	4.6	1	1
2011	5	1	2.2	2	0
2012	3	0	1.4	0	0
2013 ^e	1	0			
10-year avg.	6.7	0.2	2.8	3.7	0.1

^a Fluorescent orange requirement during fall season began fall 1992.

^b Fluorescent orange requirement for spring season began spring 1993 while moving.

^c No participation data available for 2004; 2003 data were used.

^d Fluorescent orange requirement for spring season rescinded.

^e Preliminary.

nia. Fatal and nonfatal reported separately.

ALL
(HRSIs/100,000 hunters)
5.7
7.4
8.8
7
6.4
16.2
14.7
2.8
2.7
2.9
5.4
4.8
5.6
4
6.5
4.5
5.7
6.9
0.9
?
1.0
2.2
0.6
1.3
1.9
1.2
1.4
1.9
